

# A note on the generalized $q$ -Euler numbers(2)

By

Kyoung-Ho PARK, Young-Hee KIM, and Taekyun KIM

**Abstract.** Recently the new  $q$ -Euler numbers and polynomials related to Frobenius-Euler numbers and polynomials are constructed by Kim (see[3]). In this paper, we study the generalized  $q$ -Euler numbers and polynomials attached to  $\chi$  related to the new  $q$ -Euler numbers and polynomials which is constructed in [3]. Finally, we will derive some interesting congruence on the generalized  $q$ -Euler numbers and polynomials attached to  $\chi$ .

**2000 Mathematics Subject Classification:** 11S80, 11B68.

**Key Words and Phrases:**  $q$ -Euler numbers,  $q$ -Euler polynomials.

## §1. Introduction

Let  $\mathbb{Z}, \mathbb{R}$  and  $\mathbb{C}$  denote the ring of integers, the field of real numbers and the complex number field. and let  $p$  be a fixed an odd prime number. Assume that  $q$  is an indeterminate in  $\mathbb{C}$  with  $q \in \mathbb{C}$  with  $|q| < 1$ . As the  $q$ -symbol  $[x]_q$ , we denote  $[x]_q = \frac{1-q^x}{1-q}$ . Recently,  $q$ -Euler polynomials are defined as

$$\frac{[2]_q}{qe^t + 1} e^{xt} = \sum_{n=0}^{\infty} E_{n,q}(x) \frac{t^n}{n!}, \text{ for } |t + \log q| < \pi, \text{ (see [3])}.$$

In the special case  $x = 0$ ,  $E_{n,q} = E_{n,q}(0)$  are call the  $n$ -th  $q$ -Euler numbers (see [3]). These  $q$ -Euler numbers and polynomials are closely relayed to Frobenius-Euler numbers and polynomials and these numbers are studied by Simsek-Cangul-Ozden, Cenkci-Kurt and Can and several authors (see [1-2, 18-26]). In this paper, we study the generalized  $q$ -Euler numbers and polynomials attached to  $\chi$  related to the  $q$ -Euler numbers and polynomials,  $E_{n,q}(x)$ , which is constructed in [3]. Finally, we will derive some interesting congruence on the generalized  $q$ -Euler numbers and polynomials attached to  $\chi$ .

## §2. Congruence for $q$ -Euler numbers and polynomials

The ordinary Euler polynomials are defined as

$$e^{xt} \frac{2}{e^t + 1} = e^{E(x)t} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}, \quad (\text{see [1-5]}),$$

where we use the technical method notation by replacing  $E^n(x)$  by  $E_n(x)$  ( $n \geq 0$ ), symbolically (see [1-2]). Let us consider the generating function of  $q$ -Euler polynomials  $E_{n,q}(x)$  as follows:

$$F_q(x, t) = \frac{[2]_q}{qe^t + 1} e^{xt} = \sum_{n=0}^{\infty} E_{n,q}(x) \frac{t^n}{n!}, \quad (1)$$

and we also note that

$$\sum_{n=0}^{\infty} E_{n,q}(x) \frac{t^n}{n!} = \frac{[2]_q}{qe^t + 1} e^{xt} = \frac{1 - (-q^{-1})}{e^t - (-q^{-1})} = \sum_{n=0}^{\infty} H_n(-q^{-1}, x) \frac{t^n}{n!},$$

where  $H_n(-q^{-1}, x)$  are called the  $n$ -th Frobenius-Euler polynomials (see [3]). From (1), we note that

$$\lim_{q \rightarrow 1} F_q(x, t) = \frac{2}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}. \quad (2)$$

By (1) and (2), we see that

$$\lim_{q \rightarrow 1} E_{n,q}(x) = E_n(x).$$

In (1), it is easy to show that

$$\sum_{n=0}^{\infty} E_{n,q}(x) \frac{t^n}{n!} = F_q(x, t) = \frac{[2]_q}{qe^t + 1} e^{xt} = \sum_{n=0}^{\infty} \left( \sum_{l=0}^n \binom{n}{l} E_{l,q} x^{n-l} \right) \frac{t^n}{n!}.$$

By comparing the coefficients on the both sides, we have

$$E_{n,q}(x) = \sum_{l=0}^n \binom{n}{l} E_{l,q} x^{n-l}, \quad \text{where } E_{l,q} \text{ are the } l\text{-th } q\text{-Euler numbers.} \quad (3)$$

Let  $\chi$  be the Dirichlet's character with conductor  $d \equiv 1 \pmod{2}$ . Then we define generating function of the generalized  $q$ -Euler numbers attached to  $\chi$ ,  $E_{n,\chi,q}$  as follows:

$$F_{q,\chi}(t) = \frac{[2]_q \sum_{l=0}^{d-1} \chi(l) q^l (-1)^l e^{lt}}{q^d e^{dt} + 1} = \sum_{n=0}^{\infty} E_{n,\chi,q} \frac{t^n}{n!}. \quad (4)$$

From (4), we note that

$$\lim_{q \rightarrow 1} F_{q,\chi}(t) = \frac{2 \sum_{a=0}^{d-1} \chi(a) (-1)^a e^{at}}{e^{dt} + 1} = \sum_{n=0}^{\infty} E_{n,\chi} \frac{t^n}{n!}, \quad (5)$$

where  $E_{n,\chi}$  are the  $n$ -th ordinary Euler numbers attached to  $\chi$ . By (4) and (5), we see that

$$\lim_{q \rightarrow 1} E_{n,\chi,q} = E_{n,\chi}.$$

From (5), we can also derive

$$\begin{aligned} \sum_{n=0}^{\infty} E_{n,\chi,q} \frac{t^n}{n!} &= F_{q,\chi}(t) = [2]_q \sum_{k=0}^{\infty} \chi(k) (-q)^k e^{kt} \\ &= \sum_{n=0}^{\infty} \left( [2]_q \sum_{k=0}^{\infty} \chi(k) (-q)^k k^n \right) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left( d^n \sum_{a=0}^{d-1} (-q)^a \chi(a) E_{n,q^d} \left( \frac{a}{d} \right) \right) \frac{t^n}{n!}. \end{aligned} \quad (6)$$

By comparing the coefficients on the both sides of (6), we have

$$E_{n,\chi,q} = [2]_q \sum_{k=0}^{\infty} \chi(k) (-q)^k k^n = d^n \sum_{a=0}^{d-1} (-q)^a \chi(a) E_{n,q^d} \left( \frac{a}{d} \right). \quad (7)$$

Finally, we define the generating function of the generalized  $q$ -Euler polynomials attached to  $\chi$ ,  $E_{n,\chi,q}(x)$  as follows:

$$F_{q,\chi}(x, t) = \sum_{n=0}^{\infty} E_{n,\chi,q}(x) \frac{t^n}{n!} = [2]_q \sum_{k=0}^{\infty} \chi(k) (-q)^k e^{(x+k)t}. \quad (8)$$

By (8), we easily see that

$$\begin{aligned} \sum_{n=0}^{\infty} E_{n,\chi,q}(x) \frac{t^n}{n!} &= F_{q,\chi}(x, t) = [2]_q \sum_{k=0}^{\infty} \chi(k) (-q)^k e^{(x+k)t} \\ &= \sum_{n=0}^{\infty} \left( [2]_q \sum_{k=0}^{\infty} \chi(k) (-q)^k (x+k)^n \right) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left( d^n \sum_{a=0}^{d-1} (-q)^a \chi(a) E_{n,q^d} \left( \frac{a+x}{d} \right) \right) \frac{t^n}{n!}. \end{aligned} \quad (9)$$

Thus, we have

$$E_{n,\chi,q}(x) = d^n \sum_{a=0}^{d-1} (-q)^a \chi(a) E_{n,q^d} \left( \frac{a+x}{d} \right) = \sum_{\ell=0}^n \binom{n}{\ell} x^{n-\ell} E_{\ell,\chi,q} = [2]_q \sum_{k=0}^{\infty} \chi(k) (-q)^k (x+k)^n. \quad (10)$$

Let  $d \in \mathbb{N}$  with  $d \equiv 1 \pmod{2}$ . Then, we see that

$$\begin{aligned} q^d F_{q,\chi}(d, t) + F_{q,\chi}(t) &= [2]_q \sum_{k=0}^{\infty} \chi(k) (-q)^k e^{(d+k)t} + [2]_q \sum_{k=0}^{\infty} \chi(k) (-q)^k e^{kt} \\ &= [2]_q \sum_{k=0}^{d-1} \chi(k) (-q)^k e^{kt}. \end{aligned} \quad (11)$$

From (11), we have

$$\sum_{n=0}^{\infty} \left( q^d E_{n,\chi,q}(d) + E_{n,\chi,q} \right) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left\{ [2]_q \sum_{k=0}^{d-1} \chi(k) (-q)^k k^n \right\} \frac{t^n}{n!}.$$

Therefore, we obtain the following theorem.

**THEOREM 1.** *For  $q \in \mathbb{C}$  with  $|q| < 1$ ,  $n \in \mathbb{Z}_+$  and  $d \in \mathbb{N}$  with  $d \equiv 1 \pmod{2}$ , we have*

$$q^d E_{n,\chi,q}(d) + E_{n,\chi,q} = [2]_q \sum_{k=0}^{d-1} \chi(k) (-q)^k k^n.$$

Let  $p$  be a positive odd integer and let  $N \in \mathbb{N}$ . Then we have

$$\begin{aligned} [2]_q \sum_{a=0}^{dp^N-1} \chi(a) (-q)^a a^n &= q^{dp^N} E_{n,\chi,q}(dp^N) + E_{n,\chi,q} \\ &= q^{dp^N} \sum_{j=0}^n \binom{n}{j} (dp^N)^j E_{n-j,\chi,q} + E_{n,\chi,q} \\ &= q^{dp^N} \sum_{j=1}^n \binom{n}{j} (dp^N)^j E_{n-j,\chi,q} + (q^{dp^N} + 1) E_{n,\chi,q} \\ &\equiv 2E_{n,\chi,q} \pmod{dp^N}, \end{aligned}$$

because  $q^{n dp^N} \equiv 1 \pmod{dp^N}$ . Therefore, we obtain the following theorem.

**THEOREM 2.** *Let  $p$  be a positive odd integer and  $q \in \mathbb{C}$  with  $|q| < 1$  and  $(q-1, dp) = 1$ . For  $d \in \mathbb{N}$  with  $d \equiv 1 \pmod{2}$ , we have*

$$[2]_q \sum_{a=0}^{dp^N-1} \chi(a) (-q)^a a^n \equiv 2E_{n,\chi,q} \pmod{dp^N}.$$

**REMARK.** *Define*

$$L_{E,q}(s, \chi|x) = [2]_q \sum_{n=0}^{\infty} \frac{(-q)^n \chi(n)}{(n+x)^s},$$

where  $s \in \mathbb{C}$ , and  $x \neq 0, -1, -2, \dots$ . For  $k \in \mathbb{Z}_+$ , we have  $L_{E,q}(-k, \chi|x) = E_{k,\chi,q}(x)$ .

**ACKNOWLEDGEMENTS.** The present research has been conducted by the research grant of the Kwangwoon University in 2009.

## REFERENCES

- [1] R. P. Agarwal, C. S. Ryoo, *Numerical Computations of the roots of the generalized twisted  $q$ -Bernoulli polynomials*, Neural parallel Sci. Comput, **15** (2007), 193-206.
- [2] I. N. Cangul, H. Odzen and Y. Simsek, *Generating Functions of the  $(h, q)$ -extension of Euler polynomials and numbers*, Acta Math. Hungar. **120** (2008), 281-299.
- [3] T. Kim, *An invariant  $p$ -adic  $q$ -integral on  $\mathbb{Z}_p$* , Appl. Math. Letters **21** (2008), 105-108.
- [4] T. Kim,  *$q$ -extension of the Euler formula and trigonometric functions*, Russ. J. Math. Phys. **14** (3) (2007), 275-278.
- [5] T. Kim, *Symmetry  $p$ -adic invariant integral on  $\mathbb{Z}_p$  for Bernoulli and Euler polynomials*, J. Difference Equ. Appl. **14** (2008), 1267-1277.
- [6] H. M. Srivastava, T. Kim, and Y. Simsek,  *$q$ -Bernoulli numbers and polynomials associated with multiple  $q$ -zeta functions and basic  $L$ -series*, Russ. J. Math. Phys. **12**(2) (2005), 241-268.
- [7] T. Kim, *Note on  $q$ -Genocchi numbers and polynomials*, Adv. Stud. Contemp. Math. **17** (2008), 9-15.
- [8] T. Kim, *The modified  $q$ -Euler numbers and polynomials*, Adv. Stud. Contemp. Math. **16** (2008), 161-170.
- [9] T. Kim, *On a  $q$ -analogue of the  $p$ -adic log gamma functions and related integrals*, J. Number Theory, **76** (1999), 320-329.
- [10] T. Kim,  *$q$ -Volkenborn Integration*, Russ. J. Math. Phys. **9** (2002), 288-299.
- [11] T. Kim,  *$q$ -Bernoulli numbers and polynomials associated with Gaussian binomial coefficients*, Russ. J. Math. Phys. **15** (2008), 51-57.
- [12] T. Kim, J. Y. Choi, and J. Y. Sug, *Extended  $q$ -Euler numbers and polynomials associated with fermionic  $p$ -adic  $q$ -integral on  $\mathbb{Z}_p$* , Russ. J. Math. Phys. **14** (2007), 160-163.
- [13] T. Kim, *On the von Staudt-Clausen's Theorem for the  $q$ -Euler numbers*, Russ. J. Math. Phys. **16** (3) (2009).
- [14] T. Kim,  *$q$ -generalized Euler numbers and polynomials*, Russ. J. Math. Phys. **13**(3)(2006), 293-298.

- [15] T. Kim, *Multiple  $p$ -adic  $L$ -function* Russ. J. Math. Phys. **13(2)** (2006), 151-157.
- [16] T. Kim, *Power series and asymptotic series associated with the  $q$ -analog of the two-variable  $p$ -adic  $L$ -function*, Russ. J. Math. Phys. **12(2)** (2005), 186-196.
- [17] T. Kim, *Analytic continuation of multiple  $q$ -zeta functions and their values at negative integers*, Russ. J. Math. Phys. **11(1)** (2004), 71-76.
- [18] T. Kim *On Euler-Barnes multiple zeta functions*, Russ.J. Math. Phys. **10(3)** (2003), 261-267.
- [19] B. A. Kupershmidt, *Reflection symmetries of  $q$ -Bernoulli polynomials*, J. Nonlinear Math. Phys. **12** (2005), 412-422.
- [20] H. Ozden, Y. Simsek, *Interpolation functions of the  $(h, q)$ -extension of twisted Euler numbers*, Comput. Math. Appl. **56** (2008), 898-908.
- [21] H. Ozden, Y. Simsek, *A new extension of  $q$ -Euler numbers and polynomials related to their interpolation functions*, Appl. Math. Lett. **21** (2008), 934-939.
- [22] M. Schork, *Ward's calculus sequence;  $q$ -calculus and their limit  $q \rightarrow -1$* , Adv. Stud. Contemp. Math. **13** (2006), 131-141.
- [23] Y. Simsek, *Generating functions of the twisted Bernoulli numbers and polynomials associated with their interpolation functions*, Adv. Stud. Contemp. Math. **16** (2008), 251-278.
- [24] Y. Simsek,  *$p$ -adic twisted  $q$ - $L$ -functions related to generalized twisted Bernoulli numbers*, Russ. J. Math. Phys. **13(3)** (2006), 340-348.
- [25] Y. Simsek, O. Yurekli, and V. Kurt, *On interpolation functions of the twisted generalized Frobenius-Euler numbers*, Adv. Stud. Contemp. Math. **15(2)** (2007), 187-194.
- [26] S. H. Rim, K. H. Park, and E.J. Moon, *On Genocchi numbers and polynomials*, Abstract and Applied Analysis, Article ID **898471**, (2008) 7 pages, doi:10.1155/2008/898471.
- [27] T. Kim, Y.H. Kim, K. W. Hwang *On the  $q$ -extensions of the Bernoulli and Euler numbers, related identities and lerch zeta function*, Proc. Jangjeon Math. Soc. **12** (2009), no. 1, 1798–1804.

Kyoung-Ho PARK

Department of Mathematics,  
Sogang University, Seoul 121-741, S. Korea

*E-mail:* sagamath@yahoo.co.kr

Young-Hee KIM

Division of General Education-Mathematics,  
Kwangwoon University, Seoul 139-701, S. Korea

*E-mail:* yhkim@kw.ac.kr

Taekyun KIM

Division of General Education-Mathematics,  
Kwangwoon University, Seoul 139-701, S. Korea

*E-mail:* tkkim@kw.ac.kr